

**CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM**  
**maintained by the**  
**CALIFORNIA DEPARTMENT OF FISH AND GAME**  
**and supported by the**  
**CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP**  
**Database Version 8.1 (2005)**

---

B396 Bendire's Thrasher *Toxostoma bendirei*  
Family: Mimidae Order: Passeriformes Class: Aves

Written by: D. Dobkin  
Reviewed by: L. Mewaldt  
Edited by: R. Duke, S. Granholm  
Updated by: CWHR Program Staff, August 2005

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

A very local spring and summer resident and breeder in flat areas of desert succulent shrub and Joshua tree habitats in Mojave Desert area. Occurs primarily in San Bernardino Co. and western Kern Co. (Grinnell and Miller 1944, Remsen 1978, Garrett and Dunn 1981). A recent study reported breeders more widespread than recorded previously, including in the Colorado Desert (England and Laudenslayer 1989). Migrants appear in California in February. Most have left breeding grounds by August, although fall and winter records occasionally reported from breeding areas, and to the north and west (England and Laudenslayer 1989).

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** Forages on ground on flat desert floor with scattered clumps of cactus, yucca, thorny scrub. In Arizona, commonly foraged in interspersed open ground and dense shrubs with cholla (Phillips et al. 1964). Feeds on caterpillars, beetles, other insects, and other invertebrates. Pecks and probes in soil and litter (Bent 1948).

**Cover:** Seeks cover in stands of thorny shrubs and cactus in flat desert areas.

**Reproduction:** Nests in cholla, yucca, paloverde, thorny shrub, or small tree. Nest usually placed 0.2 to 6 m (0.5 to 20 ft) above ground, averaging about 1.5 m (5 ft).

**Water:** No information found, but lives in desert.

**Pattern:** Frequents flat desert areas with scattered stands of thorny shrubs and cactus for cover, foraging, and nesting.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Yearlong, diurnal activity.

**Seasonal Movements/Migration:** Mostly migrates to southern Arizona, southwestern New Mexico, or Mexico to winter.

**Home Range:** No information found. Emlen (1974) reported density in an Arizona desert area of 0.2 per 40 ha (100 ac).

**Territory:** No information found.

**Reproduction:** Breeds from late February into early August. Clutch size 3 or 4, rarely 5.

May raise 2 broods, occasionally 3, per season (Bent 1948).

Niche: Little life history information available. A California Species of Special Concern. May be fewer than 200 pairs remaining in California population (Remsen 1978). Potentially serious threats to this species include harvesting of Joshua tree and other yuccas, grazing by domestic livestock, urbanization, and off-road vehicle activity within its limited breeding range (Emlen 1974, Remsen 1978).

## REFERENCES

- Bent, A. C. 1948. Life histories of North American nuthatches, wrens, thrashers, and their allies. U.S. Natl. Mus. Bull. 195. 475pp.
- Emlen, J. T., Jr. 1974. An urban bird community in Tucson, Arizona: derivation, structure, regulation. *Condor* 76:184-197.
- Engels, W. L. 1940. Structural adaptations in thrashers (Mimidae: Genus *Toxostoma*) with comments on interspecific relationships. *Univ. Calif. Publ. Zool.* 42:341-400.
- England, A. S., and W. L. Laudenslayer, Jr. 1989. Distribution and seasonal movements of Bendire's Thrasher in California. *West. Birds* 20:97-123.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. *Pac. Coast Avifauna* No. 27. 608pp.
- McCaskie, G., R. Stallcup, and P. De Benedictis. 1967. The status of certain fringillids in California. *Condor* 69:426-429.
- Phillips, A., J. Marshall, and G. Monson. 1964. The birds of Arizona. Univ. Arizona Press, Tucson. 220pp.
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. Calif. Dep. Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1. 54pp.